arXiv:0905.2460v1 [cond-mat.mes-hall] 15 May 2009

'Measurement-free' quantum feedback control of nuclear field squeezing in quantum dot

Wang Yao
Department of Physics, and Center for Theoretical and Computational Physics, The University of Hong Kong, Hong Kong, China

In the coupled electron-nuclear spin dynamics, the nuclear hyperfine field can perturb the spin orientation of the electron eigenstates in a moderate external field, which in turn determines the back-action to the nuclear spins. This makes possible 'measurement-free' quantum feedback control of nuclear field squeezing for enhancing electron spin dephasing time, and exemplary realizations are demonstrated using cyclic controls in a double quantum-dot system. In particular, we show that the remarkable enhancement of two-spin dephasing time observed by Reilly *et al.* [Science **321**, 817 (2008)] is a manifestation of this feedback squeezing of the nuclear field gradient between the two dots, and our theory agrees quantitatively with that experiment.

PACS numbers: 73.21.La, 03.67.Pp, 76.70.Fz, 71.70.Jp

A single electron spin in a semiconductor quantum dot is an attractive candidate for a solid-state quantum bit [1]. Recent experiments have demonstrated the capability of read, write, and control of single spin in different III-V quantum dot systems, by optical means [2, 3, 4, 5, 6, 7], or electric means [8, 9, 10, 11, 12, 13, 14]. An outstanding bottleneck towards spin-based quantum computation has been the fast dephasing of the electron spin by the nuclear spin environment in the III-V materials. Even at temperature as low as ~ 100 mK, the thermal statistical fluctuation of the nuclear spin configurations still corresponds to a large inhomogeneous broadening in nuclear magnetic field, which dephases the electron spin in a timescale of $T_2^* \sim 1-10$ ns [10, 11, 15, 16, 17]. Spin echo approach can transiently remove inhomogeneous dephasing at certain spin-echo times for an ultrashort duration equal to T_2^* time [12, 14], but this may be insufficient to allow the general quantum logic controls.

In principle, the nuclear field inhomogeneous broadening can be squeezed below its thermal value after proper procedures of nuclear state preparation [18, 19, 20, 21, 22, 23, 24], and the resultant enhancement on the T_2^* time can last for seconds or even longer as nuclear spin relaxation is extremely slow. This could be a solution for preparing a spin qubit with desirable coherence properties. For optically controllable electron spin in self-assembled dot, enhancement of T_2^* up to microsecond by nuclear state preparation has been achieved respectively for a spin ensemble [25], and for a single spin [26]. For electrically controllable spin qubit, a remarkable experiment on nuclear state preparation was also reported [27]. In an electrostatically defined double-dot system, a cyclic experimental control of the two-electron state through the resonance between the singlet and one triplet state, designed for dynamical nuclear polarization (DNP), has surprisingly enhanced the inhomogeneous dephasing time of the two-spin states to microsecond [27].

In this work, we reveal a feedback loop that naturally lies in the electron-nuclear spin dynamics: the nuclear hyperfine field can perturb the spin orientation of the electron eigenstates in a moderate external field, which in turn determines the back-action to the nuclear spin configuration. This makes possible measurement-free quantum feedback control of nuclear field squeezing for enhancing electron spin dephasing time. We show that the cyclic experimental controls adopted for DNP in coupled double-dot [27, 28, 29] can be an exemplary realization of such feedback. In particular, the enhancement in two-spin dephasing time observed in Ref. 27 is shown to be a manifestation of this feedback squeezing of the nuclear field gradient between the two dots, and our theory is in *quantitative* agreement with the experimental observations. The feedback effect on the nuclear field is shown to depend on the sign of the hyperfine constant. For the similar system of Ge/Si double quantum dot recently realized with high quality and controllability [30, 31], the same control as in Ref. 27 will magnify the fluctuations in the nuclear field gradient. We show that feedback squeezing in such system with negative hyperfine constant can also be realized by arranging the DNP cycle to pass through a different singlet-triplet resonance.

Cyclic DNP control in double-dot.—We first briefly explain the DNP cycles controlled by voltages applied to electrostatic gates in the double dot system [27, 28, 29]. A schematic energy-level diagram with (n, m) indicating the charge occupancies of the left and right dots is shown in Fig. 1. A total of five two-electron states are involved: the singlet and triplet states of the $(1, 1)$ charge configuration, and the singlet state of the $(2, 0)$ charge configuration. All other states are well detuned and can be neglected. The detuning ε between the $(2, 0)$ and $(1, 1)$ charge configurations is gate-controlled. The three triplet states (T_-, T_0, T_+) of the $(1, 1)$ configuration are split in a magnetic field. In the vicinity of the $(2, 0) - (1, 1)$ charge degeneracy, inter-dot tunneling causes the anti-crossing of the two singlet states $(2, 0)S$ and $(1, 1)S$. Away from the $(2, 0) - (1, 1)$ charge degeneracy point, the residue inter-dot tunneling results in an exchange splitting between $(1, 1)S$ and $(1, 1)T_0$, which allows the $(1, 1)S$ state to be degenerate with $(1, 1)T_+$ or $(1, 1)T_-$ at negative or positive ε. DNP are highly efficient at the $S - T_+$ and $S - T_-$ degeneracies since energy conservation can be directly satisfied [28].

In the experiment of Ref. 27, the double dot is initialized on the $(2, 0)S$ state when ε is large and positive (see Fig. 1). The two-electron wavefunction is adiabatically evolved to $(1, 1)S$ state as ε is tuned to negative by gate-control. The evolution is slowly passed through the $S - T_+$ degeneracy, where the $S \rightarrow T_+$ transition is expected to occur with a simultaneous flip down of one nuclear spin. On the state $(1, 1)T_+$, the gate-control will be rapidly brought back to the large and positive ε, waiting for the double dot to be initialized again to the $(2,0)S$ state. The cycle is iterated at the rate of 4 MHz in an external field $B_0 = 10$ mT. Measurement indicates that the DNP cycles establish a steady-state nuclear field of ~ 10 mT, corresponding to a modest nuclear polarization of $\sim 1\%$. Surprisingly, the experiment shows that the dephasing time between the spin states $|\uparrow\rangle_L|\downarrow\rangle_R$ and $|\downarrow\rangle_L|\uparrow\rangle_R$ is increased from 15 ns to ~ 1 μs after DNP pumping for ~ 1 s. Here and hereafter, $|\uparrow\rangle$ and $|\downarrow\rangle$ denote the eigenstates of $\hat{S}^z$, the component of electron spin along the external field in z-direction, and L and R denote left and right dots respectively.

Each electron is coupled to the nuclear spins in the same dot by the contact hyperfine interaction $H_{\mathrm{e-n}} = \sum_n a_n \hat{S}^z \hat{I}^z_n + \frac{1}{2} a_n (\hat{S}^+ \hat{I}^-_n + \hat{S}^- \hat{I}^+_n)$. The electron-nuclear flip-flop by $\hat{S}^+\hat{I}^-_n$ or $\hat{S}^-\hat{I}^+_n$ is responsible for the DNP when $(1,1)S$ is in resonance with $(1,1)T_+$ or $(1,1)T_-$, but is suppressed when their detuning is large. The longitudinal part $H^z_{\mathrm{e-n}} = \hat{S}^z \sum_n a_n \hat{I}^z_n$ contributes the nuclear field along the external field direction. The evolution of the relative phase between the two-spin states $|\uparrow\rangle_L|\downarrow\rangle_R$ and $|\downarrow\rangle_L|\uparrow\rangle_R$ in an arbitrary nuclear spin-configuration $|\mathcal{J}\rangle$ is then determined by the nuclear field gradient $A^z_L - A^z_R$, where $A^z_{L(R)} \equiv \langle \mathcal{J}| \sum_{n \in L(R)} a_n \hat{I}^z_n |\mathcal{J}\rangle$ is the nuclear field in the left (right) dot. Since the experiment in Ref. 27 is performed at a temperature (~ 100 mK) many orders larger than the nuclear Zeeman energy (~ 0.01 mK), the initial nuclear spin bath under thermal equilibrium has equal probability on every possible spin-configuration. In the ensemble averaged evolution, the phase coherence between $|\uparrow\rangle_L|\downarrow\rangle_R$ and $|\downarrow\rangle_L|\uparrow\rangle_R$ is determined by the ensemble averaged factor $\sum_{\mathcal{J}} e^{-i[A^z_L(\mathcal{J}) - A^z_R(\mathcal{J})]t}$. Thus, the statistical fluctuation of $A^z_L - A^z_R$ in the various possible nuclear spin configurations leads to the inhomogeneous dephasing. The observed dephasing time of 15 ns before the DNP pump is in agreement with the calculated fluctuations of a thermally distributed nuclear spin bath [27]. The enhancement in dephasing time is clearly due to the squeezing of fluctuation in the nuclear field gradient.

Feedback squeezing of the nuclear field gradient.—We show below that the nuclear field gradient perturbs the spin orientation of the singlet state in the moderate external field, inducing an electron spin gradient between the two dots. Therefore, in each DNP cycle, the flip-down of a nuclear spin is likely to occur in the left or the right dot depending on the sign of $A^z_L - A^z_R$. This feedback mechanism is the key to the squeezing of the fluctuations in the nuclear field gradient.

With a nonzero nuclear field gradient, $H^z_{\mathrm{e-n}}$ will couple $(1,1)S$ to the $(1,1)T_0$ state. Near the $(1,1)S-(1,1)T_+$ resonance, considering this off-resonance coupling, the wavefunction of the $(1,1)S$ state (labeled as ② in Fig.1) is corrected to:

$$\alpha_{+,-}|\uparrow\rangle_L|\downarrow\rangle_R - \alpha_{-,+}|\downarrow\rangle_L|\uparrow\rangle_R = \frac{|\uparrow\rangle_L|\downarrow\rangle_R - |\downarrow\rangle_L|\uparrow\rangle_R}{\sqrt{2}} - \frac{A^z_L - A^z_R}{2g\mu_B B}\frac{|\uparrow\rangle_L|\downarrow\rangle_R + |\downarrow\rangle_L|\uparrow\rangle_R}{\sqrt{2}} \qquad (1)$$

where $g\mu_B B$ is the electron Zeeman energy in the total magnetic field. If $A^z_R < A^z_L$, $|\alpha_{+,-}|^2 < |\alpha_{-,+}|^2$ so the electron spin polarization is negative in the left dot and positive in the right dot in the $(1,1)S$ configuration. Therefore, upon the $S \rightarrow T_+$ transition, the flip down of a nuclear spin is more likely to occur in the *left* dot, which decreases A^z_L and reduces $|A^z_L - A^z_R|$. If $A^z_R > A^z_L$, the electron-nuclear flip-flop is more likely to occur in the *right* dot which also reduces $|A^z_L - A^z_R|$. Thus, in a single event of electro-nuclear flip-flop, the expectation value of $|A^z_L - A^z_R|$ is unconditionally reduced by $\bar{a}\,\big||\alpha_{-,+}|^2 - |\alpha_{+,-}|^2\big| = \bar{a}\frac{|A^z_L - A^z_R|}{g\mu_B B}$, where $\bar{a}$ is the typical coupling strength between the electron spin and one nuclear spin. $\bar{a} \sim \frac{\mathcal{A}}{N} \sim 10^4$ s^{-1}, with $\mathcal{A} \sim 10^{11}$ s^{-1} being the hyperfine constant and $N \sim 10^7$ being the total number of nuclear spins [27].

For an *arbitrary* initial nuclear configuration, the evolution of $A^z_L - A^z_R$ by the DNP cycles is universally described by the Langevin equation

$$\frac{d}{dt}(A^z_L - A^z_R) = \frac{-\Gamma\bar{a}}{g\mu_B B}(A^z_L - A^z_R) + F_0(t) + F_S(t), \qquad (2)$$

where Γ is the number of nuclear spins flipped per second. In each DNP cycle, the reduction in expectation value of $|A^z_L - A^z_R|$ is the average effect of two possible events (i.e. a nuclear spin flip in dot L or in dot R), so this control has an intrinsic stochasticity described by $F_0(t) \equiv \sum_s \sigma_s \bar{a}\delta(t - s\Gamma^{-1})$, where σ_s is randomly selected from 1 and -1. $F_S(t)$ is a stochastic force describing the spectral diffusion of the nuclear field [32, 33, 34, 35, 36]. Clearly, this feedback control is only determined by the nuclear spin flip rate Γ and is *independent* of whether the electron-nuclear flip-flop process is adiabatic or not [23, 27], showing the robustness of the feedback loop. *Regardless* of the initial nuclear spin configuration, the feedback control always drives the nuclear spin bath towards a stable configuration with $A^z_L = A^z_R$.

Comparison with experiment.— The residue statistical fluctuation in the steady-state is determined by the competition of the feedback squeezing with the spectral diffusion process and the intrinsic stochasticity. Spectral diffusion in a nuclear spin bath is mainly caused by pair-flips between neighboring nuclear spins [32, 33, 34, 35, 36]. The nuclear field shift by a single pair-flip is $\sim \mathcal{A}N^{-\frac{4}{3}}$, and the number of pair-flips in a time period Γ^{-1} is $\sim Nb^2\Gamma^{-2}$, with $b \sim 10^2$ s^{-1} being the energy scale of nuclear dipolar interaction [34, 35, 36].


Energy
(1,1)T_-
⑦
(1,1)T_-
⑤
(1,1)S
⑥
(1,1)T_+
④
② (1,1)S
③ (1,1)T_+
①
(2,0)S
ε
Feedback control via $S \to T_+$
①
④
DNP in dot L
$\langle S_L^z \rangle_S < \langle S_R^z \rangle_S$ if $A_L^z > A_R^z$
$|(1,1)S\rangle$
$|(1,1)T_+\rangle$
②
③
DNP in dot R
$\langle S_L^z \rangle_S > \langle S_R^z \rangle_S$ if $A_L^z < A_R^z$

FIG. 1: Top: energy-level diagram of the coupled double-dot. The detuning ε between the $(2,0)$ and $(1,1)$ charge configurations is gate-controlled. The anti-crossing of the $(2,0)S$ and $(1,1)S$ is due to inter-dot tunneling which can be independently controlled. Bottom: feedback loop in the DNP cycle passing through the $S - T_+$ resonance in an external field along z-direction. The statistical broadening in the nuclear field gradient $A_L^z - A_R^z$ is squeezed (magnified) if the hyperfine constant $\mathcal{A}$ is positive (negative).

Thus the residue statistical fluctuation due to spectral diffusion is $\Delta(A_L^z - A_R^z) \sim \frac{b^2\Gamma^{-2}\mathcal{A}N^{-1/3}}{\bar{a}(g\mu_B B)^{-1}}$. In the experiment of Ref. 27, $B \sim 10 - 20$ mT and $\Gamma \sim 0.4 - 4$ MHz assuming each cycle can flip down a nuclear spin with a probability of $10\% - 100\%$, so $\Delta(A_L^z - A_R^z) \sim 10^4 - 10^6$ s^{-1}. The residue fluctuation due to the intrinsic stochasticity F_0 can be easily calculated: $\Delta(A_L^z - A_R^z) = \sqrt{\bar{a} g\mu_B B} \approx 10^6$ s^{-1}. Besides, the intrinsic spin dephasing has a timescale $T_2 \sim \mu$s which is not affected by this feedback control [32, 33, 34, 35, 36]. Thus, Eq. (2) predicts that the enhancement in ensemble spin dephasing time saturates at $T_2^* \sim T_2 \sim \mu$s, consistent with the finding in Ref. 27.

Eq. (2) also predicts that the statistical broadening decreases in an exponential form $\exp(-\frac{\bar{a}}{g\mu_B B}\Gamma t)$. For DNP rate $\Gamma \sim 0.4 - 4$ MHz, a DNP pump time of $1 - 0.1$ s is needed to squeeze the initial thermal fluctuation by a factor of ~ 70 down to 10^6 s^{-1}. This is consistent with the experimental observation that T_2^* is enhanced to microsecond after a DNP pump time of $\sim$ s [27]. To achieve the squeezing factor of 70, the total number of nuclear spins flipped down by the DNP cycles is $N\frac{g\mu_B B}{\mathcal{A}}\ln(70) \sim 0.01N$, exactly corresponding to the nuclear polarization of $\sim 1\%$ seen in the experiment [27].

It is obvious from Eq. (2) that the feedback depends on the sign of the hyperfine constant. As the hyperfine constants for the isotopes of Ga, As, and Al are all positive [37], the $S \to T_+$ control loop realizes negative feedback that squeezes the nuclear field gradient. In the similar system of Ge/Si double quantum dot [30, 31], since ^{29}Si and ^{73}Ge both have negative hyperfine constant [38], the same control shall result in positive feedback that magnifies the fluctuations in the nuclear field gradient.

Feedback control via $S \to T_-$ transition.— We have analyzed so far the feedback loop that lies in the DNP cycle involving the $S \to T_+$ transition. The singlet state $(1,1)S$ can also be brought into resonance with the other triplet state $(1,1)T_-$ at positive detuning ε (see Fig. 1), and we may utilize this transition as well for control of the nuclear spins.

Near the $S - T_-$ resonance, the wavefunction of the $(1,1)S$ state (labeled as ⑥ in Fig. 1) is given by,

$$\frac{|\uparrow\rangle_L|\downarrow\rangle_R - |\downarrow\rangle_L|\uparrow\rangle_R}{\sqrt{2}} + \frac{A_L^z - A_R^z}{2g\mu_B B}\frac{|\uparrow\rangle_L|\downarrow\rangle_R + |\downarrow\rangle_L|\uparrow\rangle_R}{\sqrt{2}}$$

In this case, if $A_R^z < A_L^z$, the singlet state has a larger probability on the $|\uparrow\rangle_L|\downarrow\rangle_R$ component, so the flip *up* of a nuclear spin is more likely to occur in the *left* dot upon the $S \to T_-$ transition. Thus, the $S \to T_-$ transition increases (decreases) $|A_L^z - A_R^z|$ if the hyperfine constant is positive (negative), just opposite to the effect of the $S \to T_+$ transition. By cyclic controls through the $S \to T_-$ transitions, feedback squeezing of the nuclear field gradient can be realized in the Ge/Si double dot where the hyperfine constant is negative [30, 31].

In the $S \to T_-$ control cycles, the two-electron state evolves in the sequence: ① → ⑤ → ⑥ → ⑦ → ① (see Fig. 1), which can be realized by an experimental control similar to the one used in Ref. 27. The ① → ⑤ evolution can be realized by sweeping the detuning ε with inter-dot tunneling switched off, since the detuning and tunneling are controlled by independent electrodes [27, 30].

Summary and outlook.— Feedback control is necessary to compensate unpredictable disturbances, or to make control possible when the initial state of the system is uncertain. In a standard feedback loop, a sensor measures the system variable being controlled, and the difference between the actual and desired value is then sent to a controller to manipulate the system to minimize this difference.

The example here demonstrates a new possibility for feedback control of microscopic and mesoscopic quantum systems. When two quantum systems are put into contact, their interaction can play multiple roles as best illustrated by this coupled electron-nuclear spin dynamics. Hyperfine interaction with nuclear spins causes decoherence of the electron spin and hence feedback control is necessary to reduce the uncertainty in the nuclear hyperfine field. Interestingly, the electron spin itself provides an ideal handle. In the feedback loop for controlling the mesoscopic nuclear spin bath, the electron *simultaneously* plays the roles of both key elements, i.e. the sensor and the controller. The difference between the actual and desired nuclear field value is instantaneously encoded into the electron spin state. The electron, manipulated in a predetermined manner, then shepherds the nuclear spin dynamics with the desired negative feedback. The feedback control loop does not involve explicit measurement steps.

The author acknowledges stimulating discussions with Xiaodong Xu, D. G. Steel, D. Gammon, and L. J. Sham.

[1] D. Loss and D. P. DiVincenzo, Phys. Rev. A **57**, 120 (1998).

[2] M. Atatüre, J. Dreiser, A. Badolato, A. Högele, K. Karrai, and A. Imamoglu, Science **312**, 551 (2006).

[3] X. Xu, Y. Wu, B. Sun, Q. Huang, J. Cheng, D. G. Steel, A. S. Bracker, D. Gammon, C. Emary, and L. J. Sham, Phys. Rev. Lett. **99**, 097401 (2007).

[4] J. Berezovsky, M. H. Mikkelsen, N. G. Stoltz, L. A. Coldren, and D. D. Awschalom, Science **320**, 349 (2008).

[5] E. D. Kim, K. Smirl, Y. Wu, A. Amo, X. Xu, D. G. Steel, A. S. Bracker, D. Gammon, and L. J. Sham, in *2008 CLEO/QELS Conference* (2008), vol. 1-9, pp. 3593–3594.

[6] D. Press, T. D. Ladd, B. Zhang, and Y. Yamamoto, Nature **456**, 218 (2008).

[7] D. Kim, S. E. Economou, S. C. Badescu, M. Scheibner, A. S. Bracker, M. Bashkansky, T. L. Reinecke, and D. Gammon, Phys. Rev. Lett. **101**, 236804 (2008).

[8] J. M. Elzerman, R. Hanson, L. H. Willems van Beveren, B. Witkamp, L. M. K. Vandersypen, and L. P. Kouwenhoven, Nature **430**, 431 (2004).

[9] M. Xiao, I. Martin, E. Yablonovitch, and H. W. Jiang, Nature **430**, 435 (2004).

[10] A. C. Johnson, J. R. Petta, J. M. Taylor, A. Yacoby, M. D. Lukin, C. M. Marcus, M. P. Hanson, and A. C. Gossard, Nature **435**, 925 (2005).

[11] F. H. L. Koppens, J. A. Folk, J. M. Elzerman, R. Hanson, L. H. W. van Beveren, I. T. Vink, H. P. Tranitz, W. Wegscheider, L. P. Kouwenhoven, and L. M. K. Vandersypen, Science **309**, 1346 (2005).

[12] J. R. Petta, A. C. Johnson, J. M. Taylor, E. A. Laird, A. Yacoby, M. D. Lukin, C. M. Marcus, M. P. Hanson, and A. C. Gossard, Science **309**, 2180 (2005).

[13] F. H. L. Koppens, C. Buizert, K. J. Tielrooij, I. T. Vink, K. C. Nowack, T. Meunier, L. P. Kouwenhoven, and L. M. K. Vandersypen, Nature **442**, 766 (2006).

[14] F. H. L. Koppens, K. C. Nowack, and L. M. K. Vanders, Phys. Rev. Lett. **100**, 236802 (2008).

[15] A. S. Bracker, E. A. Stinaff, D. Gammon, M. E. Ware, J. G. Tischler, A. Shabaev, A. L. Efros, D. Park, D. Gershoni, V. L. Korenev, et al., Phys. Rev. Lett. **94**, 047402 (2005).

[16] M. V. G. Dutt, J. Cheng, B. Li, X. Xu, X. Li, P. R. Berman, D. G. Steel, A. S. Bracker, D. Gammon, S. E. Economou, et al., Phys. Rev. Lett. **94**, 227403 (2005).

[17] I. A. Merkulov, A. L. Efros, and M. Rosen, PRB **65**, 205309 (2002).

[18] D. Stepanenko, G. Burkard, G. Giedke, and A. Imamoglu, Phys. Rev. Lett. **96**, 136401 (2006).

[19] G. Giedke, J. M. Taylor, D. D'Alessandro, M. D. Lukin, and A. Imamoglu, Phys. Rev. A **74**, 032316 (2006).

[20] D. Klauser, W. A. Coish, and D. Loss, Phys. Rev. B **73**, 205302 (2006).

[21] G. Ramon and X. Hu, Phys. Rev. B **75**, 161301 (2007).

[22] M. S. Rudner and L. S. Levitov, Phys. Rev. Lett. **99**, 036602 (2007).

[23] H. Ribeiro and G. Burkard, arXiv:0811.3560 (2008).

[24] I. T. Vink, K. Nowack, F. Koppens, J. Danon, Y. Nazarov, and L. Vandersypen, arXiv:0902.2659 (2009).

[25] A. Greilich, A. Shabaev, D. R. Yakovlev, A. L. Efros, I. A. Yugova, D. Reuter, A. Wieck, and M. Bayer, Science **317**, 1896 (2007).

[26] X. Xu, W. Yao, B. Sun, D. G. Steel, A. S. Bracker, D. Gammon, and L. J. Sham, submitted (2008).

[27] D. J. Reilly, J. M. Taylor, J. R. Petta, C. M. Marcus, M. P. Hanson, and A. C. Gossard, Science **321**, 817 (2008).

[28] J. R. Petta, J. M. Taylor, A. C. Johnson, A. Yacoby, M. D. Lukin, C. M. Marcus, M. P. Hanson, and A. C. Gossard, Phys. Rev. Lett. **100**, 067601 (2008).

[29] S. Foletti, J. Martin, M. Dolev, D. Mahalu, V. Umansky, and A. Yacoby, arXiv:0801.3613 (2008).

[30] Y. Hu, H. O. H. Churchill, D. J. Reilly, J. Xiang, C. M. Lieber, and C. M. Marcus, Nature Nanotechnology **2**, 622 (2007).

[31] C. B. Simmons, B. M. Thalakulam, M. Rosemeyer, B. J. Van Bael, E. K. Sackmann, D. E. Savage, M. G. Lagally, R. Joynt, M. Friesen, S. N. Coppersmith, and M. A. Eriksson, arXiv:0905.1647 (2009).

[32] R. de Sousa and S. Das Sarma, Phys. Rev. B **68**, 115322 (2003).

[33] W. M. Witzel, R. de Sousa, and S. Das Sarma, Phys. Rev. B **72**, 161306(R) (2005).

[34] W. Yao, R. B. Liu, and L. J. Sham, Phys. Rev. B **74**, 195301 (2006).

[35] R. B. Liu, W. Yao, and L. J. Sham, New J. Phys. **9**, 226 (2007).

[36] W. Yao, R. B. Liu, and L. J. Sham, Phys. Rev. Lett. **98**, 077602 (2007).

[37] D. Paget, G. Lampel, and B. Sapoval, Phys. Rev. B **15**, 5780 (1977).

[38] R. K. Harris, in *Encyclopedia of Nuclear Magnetic Resonance*, edited by D. M. Granty and R. K. Harris (John Wiley & Sons, Chichester, UK, 1996), vol. 5.